

**MANGO TREE NAMED 'B74'**

5      LATIN NAME OF THE GENUS AND SPECIES OF THE PLANT CLAIMED

*Mangifera indica*

**VARIETY DENOMINATION**

**'B74'**

10

**BACKGROUND AND SUMMARY OF THE INVENTION**

15      This invention relates to the discovery and asexual propagation of a new variety of mango tree, as herein described and illustrated. The new variety was first hybridized by controlled pollination. The new variety is a precocious, heavy-cropping, upright tree yielding red-skinned, medium-sized, terpinolene-flavored fruit.

    The seed parent is 'Sensation' and the pollen parent is 'Kensington Pride.' The new variety was selected and evaluated at the fruiting stage on the property of Mr. And Mrs. L. W. Dorrian at Childers, in Queensland, Australia.

20      The new mango tree variety was first asexually propagated by grafting onto seedling rootstocks in Childers, in Queensland, Australia.

    The new mango tree variety cv. 'B74' has been shown to maintain its distinguishing characteristics through successive asexual propagations.

**BRIEF DESCRIPTION OF THE PHOTOGRAPHS**

25      The accompanying photographic illustrations illustrate in full color the new mango variety 'B74.' The colors are as nearly true as is reasonably possible in a color representation of this type.

    FIG. 1 is a photograph of the fruit of the new mango variety 'B74' (center) and comparators 'Kensington Pride' (bottom) and 'R2E2' (top) showing differences in size and color.

30      FIG. 2 is a photograph of the tree of the new mango variety 'B74.'

FIG. 3 is a photograph of the floral panicle of the new mango variety 'B74.'

FIG. 4 is a photograph of flowers and floral buds of the new mango variety 'B74.'

FIG. 5 is a photograph of a single flower of the new mango variety 'B74.'

5 FIG. 6 is a photograph of a cross-sectional slice of the fruit of the new variety 'B74' sliced through the fruit in the region above the seed.

FIG. 7 is a photograph of the seed of the new mango variety 'B74.'

#### DETAILED BOTANICAL DESCRIPTION OF THE INVENTION

Throughout this specification, color names beginning with a small letter signify  
10 that the name of that color, as used in common speech, is aptly descriptive.

The descriptive matter which follows pertains to 'B74' mango trees (as well as the comparative varieties 'Sensation,' 'Kensington Pride', and 'R2E2', each of the foregoing which is unpatented) grown in the vicinity of Childers, Queensland, Australia. The scions of the candidate and comparator varieties were topworked to  
15 'Keitt' trees (unpatented) that were originally grafted to polyembryonic seedlings of 'Kensington Pride.' Ten single tree replicates of each cultivar were planted at 6 x 10 m intervals in red basaltic soil (kraznozem) following a completely randomised design. Pest and disease treatments were applied as required. Irrigation and fertilizer application followed commercial practice. 10-20 random measurements of each  
20 characteristic were obtained from each replicate. Redness of skin color was determined using a Minolta Chroma Meter CR-200 to measure the hue angle (H). Mean values were taken from measurements at three points from the shoulder to the basal end of the sun-exposed side of each fruit. The lower the hue angle, the greater the red coloration. The observations described herein are believed to apply to plants of the variety grown  
25 under similar conditions of soil and climate elsewhere.

Ten single tree replicates of each cultivar were planted at 6 x 10 m intervals following a completely randomised design. The trees were grown on a red basaltic clay loam (kraznozem) near Childers in south east Queensland, Australia (latitude 25°S, altitude 40 m). Irrigation was available to supplement the average rainfall of 900 mm.  
30 Daily light interception (measured as photosynthetic photon flux) by tree canopies

during the period of fruit development (October to February) ranges from 15.5 to 59.5 mol quanta m<sup>-2</sup> at this site. The mean annual maximum/minimum temperatures at this site are 24/15°C (Australian Bureau of Meteorology).

5 Pest and disease treatments were applied as required. Irrigation and fertilizer application followed commercial practice. 10 to 20 random measurements of each characteristic were obtained from each replicate.

Redness of skin color was determined using a Minolta Chroma Meter CR-200 to measure the hue angle (H). Mean values were taken from measurements at three points from the shoulder to the basal end of the sun-exposed side of each fruit. The  
10 lower the hue angle, the greater the red coloration. Elsewhere throughout the specification, color references are made to the Royal Horticultural Society Color Chart. The observations described herein are believed to apply to plants of the variety grown under similar conditions of soil and climate elsewhere.

The new mango tree cultivar, 'B74' is quite distinct from its seed parent  
15 'Sensation', and may be distinguished from its seed parent in the following characteristics: The average fruit weight of 'B74' (457 g) is larger than 'Sensation' (360 g). Additionally, the new mango tree cultivar matures 3-4 weeks earlier than the seed parent 'Sensation', which is a very late variety. The skin color of the new mango tree cultivar is red and yellow, while the skin color of the seed parent 'Sensation' has a  
20 bright yellow background with a dark red to purple blush that covers most of the surface. Further, while the new mango tree cultivar has a terpinolene aroma in both the leaves and fruit, the seed parent 'Sensation' has no distinguishable terpinolene aroma in either leaves or fruit.

The new mango tree cultivar may be distinguished from the pollen parent,  
25 'Kensington Pride,' by the following characteristics. The tree of the new mango cultivar 'B74' has a more erect form and lower vigor than the tree of the pollen parent 'Kensington Pride'. The fruit of the new mango tree cultivar matures in late season, while the fruit of 'Kensington Pride' matures early to mid-season. The predominant skin color of ripe fruit of the new mango tree cultivar is approximately equal amounts  
30 of yellow and red blush (approximately 30% to 55% red blush), while the predominant

skin color of ripe fruit of 'Kensington Pride' is yellow with a small proportion of red blush. The fruit flesh of the new mango tree cultivar is pale yellow in color, while the fruit flesh of 'Kensington Pride' is yellow. The fruit shape of the new mango tree cultivar is broad elliptic, while fruit shape of the pollen parent 'Kensington Pride' is medium elliptic. The sinus proximal of the styler scar of the new mango tree cultivar is absent, while it is present in the pollen parent 'Kensington Pride.' The fruit of the new mango tree cultivar is somewhat smaller than that of 'Kensington Pride' (457 g as compared to 475 g). The seed of the new mango tree cultivar is of the monoembryonic type, while the seed of the pollen parent 'Kensington Pride' is polyembryonic.

The new mango tree variety 'B74' may be distinguished from presently available cultivars. The new mango tree cultivar 'B74' can be compared, for example, to 'R2E2', which is commonly grown in Australia, by the following distinguishing characteristics: The leaves of the new mango tree variety are shorter and wider than the leaves of 'R2E2.' The new mango tree cultivar has a higher percentage of bunch bearing inflorescences than 'R2E2.' The fruit of 'B74' is smaller than that of 'R2E2.' The flesh color of the new mango tree variety is pale yellow, while the flesh color of 'R2E2' is yellow. The skin color of ripe fruit of the new mango tree variety is approximately equal amounts of yellow and red blush (approximately 30% to 55% red blush), while the predominant skin color of ripe fruit of 'R2E2' is predominantly yellow with a small proportion of red blush. Additional comparative information can be found in Table 1.

## TREE

### General:

*Vigor:* low to moderate. Fruiting B74 mango trees growing at the experimental site typically produce only one flush of shoot growth annually following the completion of fruit harvest in early February. Autumn/winter temperatures typically restrict further vegetative growth prior to flowering in September-October. The mean length of summer shoot growth is approximately  $26.5 \pm 2.4$  mm ( $\pm$  values are the standard error of the mean).

5                    *Height and spread of tree:* At the time the description was made for Plant Breeders Rights in Australia the trees were 3 m high with a spread of 3.5 m. However, these trees were top-worked to existing 6-year-old trees. More typically nursery-produced 'B74' trees grafted to 'Kensington Pride' seedling rootstock reach approximately 1.25 m diameter and approximately 1.5 m in height by their third flowering anniversary after planting which is their first year of commercial cropping at Childers in SE Queensland.

10                   *Density of foliage:* open.

10                   *Shape:* upright.

*Form:* erect.

15                   *Fruit bearing:* mid-late season maturity. Fruit maturity of 'B74' is generally reached after the accumulation of 1650 degree days (measured by the number of °C hours accumulated above 10°C starting from the point of full inflorescence extension). At the site where the description for 'B74' was developed, fruit reached the mature green stage during the first week of February (midseason).

*Root stock:* 'Kensington Pride'.

**Trunk:**

20                   The top-worked trees used for the original description for Plant Breeders Rights in Australia did not have 'B74' trunks, as they were top-worked onto 'Keitt' trees as described above. However, the trunk data supplied below have been collected from another block of 10-year-old 'B74' trees in the same orchard where 'B74' was grafted directly to seedling 'Kensington Pride' rootstocks. The position where the diameter was

25                   measured is above the graft union.

*Mean diameter of trees measured 50 cms above ground level:* approximately 19.3 ± 0.3 cm.

*Trunk color:* about Grey-brown 199A.

30                   *Bark texture:* slightly rough with slight vertical ridges 10 to 20 mm apart.

## BRANCHES

The branch values listed below were measured on fully mature 10-year-old 'B74' trees grafted to 'Kensington Pride' seedling rootstock.

*Branch diameter at the base:* approximately  $10.08 \pm 0.24$  cm.

5 *Crotch angle:* varies from 45 to 80°.

*Branch color:* about Grey-brown 199A.

*Lenticels:* not visible.

*Surface texture:* slightly rough with slight vertical ridges 10 to 20 mm apart.

## LEAVES

### 10 **Young Leaf:**

*Anthocyanin:* present.

*Anthocyanin location on leaves:* leaf lamina.

*Anthocyanin coloration:* about Greyed-orange 164A.

*Intensity of color:* strong.

### 15 **Fully Expanded Leaf:**

*Average length:* approximately 204 mm.

*Average width:* approximately 62 mm.

*Length/width ratio:* low (approximately 3.5).

*Attitude:* horizontal.

20 *Leaf surface:* smooth.

*Profile:* concave cross section.

*Shape:* elliptic.

*Cross-section shape:* concave.

*Leaf blade tip:* acuminate.

25 *Base:* acute.

*Symmetry:* asymmetric.

*Undulation of margin:* not undulated.

*Curvature of midrib:* present.

*Position of midrib curvature:* apical.

30 *Twisting of blade:* absent.



*Inflorescence rachis color:* about Red 47C.

*Inflorescence texture:* the rachis of the inflorescence moderate pubescence.

*Mean number of inflorescences per tree:* approximately  $316.8 \pm 27.2$ .

*Mean number of flowers per inflorescence:* approximately  $764.0 \pm 32.7$ .

- 5      *Date of first and full blooms:* Variable. Flowering dates vary between years and are typically influenced by the size of the preceding crop, the time that floral induction temperatures occur in the autumn/early winter and the temperatures received during the latter part of winter. Typically at the Childers site inflorescence bud break occurs in mid to late July but
- 10      full bloom does not occur until the second to third week in September. To calculate Heat Accumulation Units to determine fruit maturity the time of full bloom in September is used.

## FLOWERS

### General:

- 15      *Diameter of open flowers:* approximately  $7.69 \pm 0.37$  mm.
- Percent hermaphrodite flowers per inflorescence:* approximately  $34.4 \pm 6.0$ .
- Percent male flowers per inflorescence:* approximately  $65.6 \pm 6.0$ .
- Floral fertility:* the flowers of 'B74' are self fertile and do not require cross pollination for fruit set.

### 20      Petals:

*Number of petals per flower:* 5.

*Petal size:*

*Length:* approximately  $3.26 \pm 0.09$  mm. *Diameter:* approximately  $2.57 \pm 0.02$  mm.

- 25      *Petal texture:* slightly pubescent.

*Petal shape:* ovate.

*Petal color:*

*Upper surface:* about Yellow-white 158D.

*Lower surface:* about Yellow-white 158C.



**Sepals:**

*Number of sepals per flower:* 5.

*Sepal size:*

*Length:* approximately  $3.78 \pm 0.09$  mm.

5 *Diameter:* approximately  $2.94 \pm 0.04$  mm.

*Sepal texture:* densely pubescent.

*Sepal shape:* ovate.

**Stamens:**

*Stamen number in hermaphrodite flowers:* 5; typically only 2 are fertile.

10 *Filament length:* approximately  $11.93 \pm 0.06$  mm

*Filament color:* about Green-white 157D.

*Anther length:* approximately  $0.86 \pm 0.06$  mm.

*Anther shape:* ovate.

*Anther color:* about Yellow-white 158B.

15 *Pollen color:* about Yellow-white 158C.

*Pollen production:* moderately prolific; pollen viability: approximately  $72.6 \pm 1.5\%$ .

**Pistils:**

*Pistil number:* 1.

20 *Pistil shape:* slightly curved.

*Pistil length:* approximately  $0.94 \pm 0.16$  mm.

*Ovary number:* 1.

*Diameter of ovary:* approximately  $1.36 \pm 0.05$  mm.

*Ovary color:* about Yellow-white 158B.

25

**FRUIT****General Harvest Characteristics:**

*General fruiting characteristics:* bunch bearing with 2-4 fruits commonly carried on each inflorescence.

30 *Bearing:* When well-managed, 'B74' is a regular cropping variety producing consistent yields each year.

*Season maturity:* mid-late season maturity.

5 *Fruit yield:* Field planted nursery-grown trees begin commercial fruiting following their 3rd flowering anniversary in the ground. For example, a nursery-grown 'B74' tree planted at Childers in the spring of 2001 will pass its first flowering anniversary in July/August 2002. The third flowering anniversary is July/August 2004 when it will set its first commercial crop. Yields recorded for the first three cropping years in a trees life are:

*Year 1:*  $11.5 \pm 1.13$  kg/tree.

10 *Year 2:*  $28.84 \pm 1.93$  kg/tree.

*Year 3:*  $37.83 \pm 1.11$  kg/tree.

At Childers orchard, spacing is 8 x 3 m (412 trees/ha). The original small orchard planted in 1993 has a fully mature canopy producing 27 t/ha in its 10th cropping year.

#### 15 **Shipping and Keeping Characteristics of Fruit:**

*Shipping and keeping quality of fruit:* 'B74' mango fruit typically reached the eating ripe stage within 9-10 days of harvest under the following ripening conditions: 10 ppm ethylene for two days at 20°C followed by holding at 20°C (no ethylene) until eating ripe stage. Once fruit reaches the eating ripe stage, it remains firm and in an acceptable saleable condition for a further 9-10 days when held at 20°C.

20 *Long term storage:* Long term storage assessments have not been completed on this variety but early indications are that it will hold in a firm condition for 18-21 days at 12°C and then ripen to an acceptable saleable condition.

25 *Bruising:* Bruising of mature green 'B74' fruit in transit has not been of commercial significance, with fruit travelling up to 3000 km by road transport.

## MATURE FRUIT

### General:

- Length:* medium - approximately 101.3 mm.  
*Width:* medium - approximately 91.3 mm.  
5 *Length/width ratio:* medium - approximately 1.1.  
*Shape:* ovate.  
*Cross-sectional shape:* broad elliptic.  
*Average weight:* approximately 457 g.  
*Stalk cavity:* shallow.  
10 *Sinus proximal of stylar scar:* absent.  
*Bulge proximal of stylar scar:* absent.

### Skin of Mature Fruit:

- Ground color:* about Yellow-orange 20A.  
*Anthocyanin coloration (blush):* skin develops high levels of red anthocyanin  
15 *pigmentation (about Red 4A) where exposed to sun.*  
*Percentage of blush covering of fruit:* approximately 30-55%.  
*Skin thickness:* thin; about 1.3 mm.  
*Adherence of flesh to skin:* moderate.  
*Lenticel spotting:* occurs over the surface of the skin and is more prominent on  
20 *the blush areas.*  
*Lenticel spot color:* about Green-yellow 1C.  
*Lenticel size:* less than approximately 0.4 mm in diameter.

## RIPE FRUIT

### Skin of Ripe Fruit:

- 25 *Redness of skin color (hue angle):* approximately 44.7 (using a Minolta Chroma Meter CR-200).

### Flesh of Ripe Fruit:

- Main color of flesh:* about Yellow-orange 15A.  
*Firmness of flesh:* firm. The firmness of ripe fruit determined by penetrometer  
30 *is approximately  $2.0 \pm 0.29$  Kg.*

*Flavor characteristics:* approximately  $14.73 \pm 1.44$  °Brix and  $0.183 \pm 0.072$  % acidity.

*Texture of flesh:* smooth.

*Amount of non-fleshy fiber in flesh attached to stone:* low.

5 *Turpinolene flavor:* present.

*Sap exudation at harvest:* present with sap burn and skin browning.

## SEED AND ENDOCARP

*Endocarp surface:* moderately fibred with the flesh adhering to its surface.

*Endocarp surface color:* about Yellow-orange 17B due to the adhering flesh.

10 *Endocarp plus seed dimensions:*

*Mean weight:* approximately  $30.92 \pm 1.42$  g.

*Mean length:* approximately  $67.67 \pm 0.23$  mm.

*Longest diameter:* approximately  $40.10 \pm 0.89$  mm.

*Monoembryonic seed removed from the endocarp:*

15 *Mean weight:* approximately  $15.27 \pm 0.66$  g.

*Mean length:* approximately  $57.26 \pm 1.15$  mm

*Longest diameter:* approximately  $31.38 \pm 0.94$  mm.

*Seed color:* about Yellow-white 158D. *Seed size:* small.

*Polyembryony:* monoembryonic.

20 The numerical data presented in the Australian Plant Breeders Rights description of 'B74' (see Table 1, below) have been derived from trees planted in a replicated, randomised block design with the mean values derived from an Analysis of Variance (Genstat 8.1). LSD = Least Significant Difference and is the value determined to differentiate statistical differences between the candidate variety and its  
25 comparators. Sig = the level at which statistical difference has been determined, i.e. P<0.01 indicates a 99% confidence level in the result obtained. "ns" = not significant and indicates that there is no statistical difference in this parameter between the candidate variety and the comparator.

30

## TABLE 1

## COMPARISON OF 'B74' WITH OTHER MANGO VARIETIES

	'B74'	'Kensington Pride'	'R2E2'
<b>YOUNG LEAF</b>			
Anthocyanin	present	present	present
Anthocyanin hue	red	red	red
Intensity of color	strong	strong	strong
<b>MATURE LEAF</b>			
Terpinolene aroma	present	present	absent
Cross-section shape	concave	concave	straight
Relief of upper surface	slightly sunken between veins	raised between veins	raised between veins
Shape of tip	acuminate	attenuate	acuminate
Shape of base	acute	acute	rounded
Predominant shape	elliptic	elliptic	elliptic
Symmetry	asymmetric	asymmetric	asymmetric
Curvature of midrib	present	present	present
Attitude	horizontal	horizontal	horizontal
<b>Petiole length (mm)</b>			
Mean	39.8	21.3	35.5
std deviation	0.7	0.7	1.0
LSD/sig	0.6	P≤0.01	P≤0.01
<b>Lamina length (mm)</b>			
Mean	203.9	181.5	236.8
std deviation	3.9	6.2	4.4
LSD/sig	2.0	P≤0.01	P≤0.01
<b>Lamina width (mm)</b>			
Mean	62.3	39.7	51.8
std deviation	0.8	1.1	0.8
LSD/sig	1.30	P≤0.01	P≤0.01
<b>Length/width ratio</b>			
Mean	3.44	4.00	4.00
std deviation	0.89	1.11	0.96
LSD/sig	0.21	P≤0.01	P≤0.01
<b>INFLORESCENCE</b>			
<b>Percentage of bunch-bearing inflorescences</b>			
Mean	66.4	22.9	34.3
std deviation	9.0	9.1	10.1
LSD/sig	11.7	P≤0.01	P≤0.01
<b>MATURE FRUIT</b>			
Cross-sectional shape	broad elliptic	medium elliptic	broad elliptic
Depth of stalk cavity	shallow	medium	medium
Sinus proximal of stylar scar	absent	present	absent
Bulge proximal of stylar scar	absent	absent	absent

Ripe fruit: predominant skin color	approximately equal amounts of yellow and red blush	predominantly yellow with a small proportion of red blush	predominantly yellow with a small proportion of red blush
Ripe fruit: predominant flesh color	pale yellow	yellow	yellow
Ripe fruit: amount of fiber in flesh	low	medium	low
attached to stone			
Ripe fruit: terpinolene flavor	present	present	absent
Length (mm)			
Mean	101.23	113.52	117.11
std deviation	2.68	2.14	3.34
LSD/sig	3.43	P≤0.01	P≤0.01
Width (mm)			
Mean	91.28	87.94	111.98
std deviation	2.80	2.71	2.43
LSD/sig	3.17	P≤0.01	P≤0.01
Length/Width ratio			
mean	1.11	1.29	1.05
std deviation	0.01	0.03	0.01
LSD/sig	0.02	P≤0.01	P≤0.01
Weight (g)			
mean	457.4	475.1	802.7
std deviation	38.1	37.0	53.0
LSD/sig	50.5	ns	P≤0.01
*Ripe color (hue angle)			
mean	44.73	67.36	53.50
std deviation	4.18	2.05	2.58
LSD/sig	3.41	P≤0.01	P≤0.01
Embryonic type	monoembryonic	Polyembryonic	mostly polyembryonic
TREE			
Form	erect	Spreading	erect
Vigor	low-moderate	high	moderate
Fruit maturity season	late	Early-mid	mid-late

\* Redness of skin color was determined using a Minolta Chroma Meter CR-200 to measure the hue angle (H).

**TABLE 2**  
**COMPARISON OF 'B74' FRUIT COLOR WITH OTHER MANGO**  
**VARIETIES**

5

Variety	Skin color	Flesh color
---------	------------	-------------

	<b>Ground color</b>	<b>Blush color</b>	
B74	20A (yellow-orange)	4A (red)	15A (yellow-orange)
Kensington Pride	20A (yellow-orange)	26B (orange)	21B (yellow-orange)
R2E2	22A (yellow-orange)	179A (greyed-red)	23A (yellow-orange)